

NGS – variant analysis

Sequencing and alignment





Image from: G. Silva (2016)

Illumina sequencing


- Sequencing-by-synthesis: 2nd generation sequencing
- Massive throughput: up to 500×10^9 bases/run
- Most used platform today

Illumina sequencing

- 50 – 300 bp
- Paired-end (or single-end)



Illumina library prep



fastq

fasta + basequality (fasta + q = fastq)

$$BASEQ = -10 \log_{10} \Pr\{\text{base is wrong}\}$$

$$-10 \log_{10} (0.01) = 20$$

$$-10 \log_{10} (0.1) = 10$$

$$-10 \log_{10} (0.5) = 3$$

fastq

reads.fastq

```
@D00283R:66:CC611ANXX:4:2311:2596:2330 1:N:0:TCCGGAG  
ACTCTACGCTCAATAAAGATTCTGATACGGCTCCTGAAATGCAGAATGAGT  
+  
B/<<<B<FFFFFFFFBBFFFFBFFFFBFFFF/FFFFFFF/BFFFFFFBFFF
```


title, starts with @

nucleotide sequence

optional description


base quality



Base quality (phred)




- S - Sanger Phred+33, raw reads typically (0, 40)
- X - Solexa Solexa+64, raw reads typically (-5, 40)
- I - Illumina 1.3+ Phred+64, raw reads typically (0, 40)
- J - Illumina 1.5+ Phred+64, raw reads typically (3, 41)
with 0=unused, 1=unused, 2=Read Segment Quality Control Indicator (**bold**)
(Note: See discussion above).
- L - Illumina 1.8+ Phred+33, raw reads typically (0, 41)

Quiz Question 3






in phase



out of phase


Illumina - limitations

- Bridge amplification
- Lengths are limited by out-of-phase of signal



Long reads (3rd generation)

- Crux: maximizing signal from a single-molecule base read-out
- Single molecule, so no out-of-phase signal
- Two frequently used platforms:
 - PacBio SMRT sequencing
 - Oxford Nanopore Technology




Long reads


- More error -> difficulties for variant analysis
- But:
 - PacBio CCS: high baseQ + no bias
 - Long reads can have higher mapping qualities
 - Long reads improve haplotyping

What to sequence?

- Whole genome/metagenome
- Reduced representation:
 - Bait capture/whole exome sequencing
 - Restriction Enzyme based (e.g. RAD seq)
 - Amplicon sequencing
 - RNA-seq

Read alignment (phred)






Software

- Basic alignment:
 - bowtie2
 - bwa-mem
- Long reads:
 - minimap2

Mapping quality



$$MAPQ = -10 \log_{10} \Pr\{\text{mapping position is wrong}\}$$

$$-10 \log_{10} (0.01) = 20$$

$$-10 \log_{10} (0.5) = 3$$

Quiz Question 4

sam

sequence alignment format



sam header

@HD VN:1.0 SO:coordinate


@SQ SN:U00096.3 LN:4641652

@PG ID:bowtie2 PN:bowtie2 VN:2.4.1 CL: bowtie2-align-s --wrapper basic-0 -x ref.fasta -1 reads_1.fastq -2 reads_2.fastq"

SAM column	example
read name	SRR519926.5
flag	89
reference	chr20
start position	61
mapping quality	42
CIGAR string	150M
reference name mate is mapped	=
start position mate	476
fragment length	515
sequence	CATCACCATCCCCAC
base quality	@>4:4C@89+&9CC@
optional	AS:i:-2
optional	XN:i:0

Quiz Question 5

Marking duplicates



Marking duplicates

- Variant calling: each read is an independent observation of the genome
- Duplicates (can) have the same molecular origin -> not independent
- Removing duplicates probably doesn't have a big effect on variant analysis

Ebbert MTW et al. (2016) Evaluating the necessity of PCR duplicate removal from next-generation sequencing data and a comparison of approaches. BMC Bioinformatics.

Unique Molecular Identifiers

- UMI added before PCR reaction
- Detect PCR duplicates and PCR errors